


## NAT Breakfast Task

Name: A.B.

Date: 19.09.2016

Set-up:



Evaluation:

Left		Right	
toast (minimum 6 slices)	✓	toaster	✓
jam with loosened screw cap	✓	butter/margarine	✓
cup	✓	1 knife	✓
teabags in a carton	✓	1 table spoon	
		1 tea spoon	
		kettle filled with water	✓
		sugar in container	
	4/4		4/7

Lateralized attention score = <sup>right</sup> contralateral - <sup>left</sup> ipsilesional proportion:  $0,57 - 1 = -0,43$

Notes:

The patient attended the right side only after repetitive request.

### Accomplishment Score:

Toast:

- toasted bread (1)
- apply butter or jam to the bread (1)
- apply butter *and* jam to the bread (2)

3/3

Tea:

- switch on the kettle (1)
- put the teabag in the cup (1)
- infuse water (1)
- add sugar (1)

2/4

Sum:

5/7

**Error Score:**

- toasts more than one slice of bread:
- mixes up the order:
  - fill in water then plug in the kettle
  - infuse water, then turn on the kettle
  - turn on the toaster without bread inside
  - butter the bread first and then put it in the toaster
  - apply jam to bread and then butter it
  - omit to stir the tea
  - omit to spread the butter and/or jam
- use one ingredient extremely excessive or sparsely:
- typical errors of apraxia:
  - spread with the finger
  - spread with the spoon
  - put the bread on top of the toaster
  - press the bread with the hand in the toaster
  - take butter with the spoon (for jam ok!)
  - take sugar with the knife
  - empty the sugar directly in the cup
  - put jam in the cup
  - put sugar on the bread
  - put the teabag on the bread

Number:

1  
1

1  
0

Error Score = Sum of errors:

3 /19

**NAT Score:**

Accomplishment Score	Error Score	Naturalistic Action Task Score	Interpretation
7	0-1	6	
7	> 1	5	no apraxia
6	0-1	4	
6	> 1	3	mild apraxia
4-5	0-1	2	moderate apraxia
4-5	> 1	1	severe apraxia
0-3	≥ 0	0	

The patient is not able to execute all single step tasks. Further, he forgot the precise instructions very fast.